

Implications of current LHC results for supersymmetry and neutralino dark matter

Sabine Kraml
(LPSC Grenoble)


Bethe Forum

Inaugural Program on LHC, Dark Matter and Unification
Bonn, Germany, 2nd – 18th November 2011

The Standard Model


<1973: theoretical foundations of the SM

- renormalizability of $SU(2) \times U(1)$ with Higgs mech. for EWSB
- asymptotic freedom, QCD as gauge theory of strong force
- KM description of CP violation



Followed by more than 30 years of consolidation

- **experimental** verification via **discovery** of
 - gauge bosons: gluon, W, Z
 - matter fermions: charm, 3rd family
- experimental **precision measurements**
 - EW radiative corrections
 - running of the strong coupling
 - CP violation in the 3rd generation
- technical **theoretical advances** (higher-order calculations...)



Only missing piece: the Higgs ?

NB: Unitarity tells us that the dynamics of EWSB should become apparent around the TeV scale.



... and beyond

The Standard Model is **tremendously successful**,
nevertheless it can't be the ultimate theory. Too many open questions.

- Why are there 3 generations of quarks and leptons?
- What is the origin of flavour mixing and CP violation?
- Why is the SM anomaly free?
- Can the different interactions (and matter fields) be unified?
- What stabilizes the electroweak scale?
-

Attempts to answer these questions, in particular regarding the naturalness and hierarchy problem, let us **expect new physics at TeV energies**.

(another) **genuine motivation to build the LHC**

Besides, the experimental evidences of **neutrino masses**, **dark matter** and the **baryon asymmetry** tell us that we are missing something fundamental.




Supersymmetry

The beauties of SUSY:



- Unique extension of space-time symmetries
- Solution to the gauge hierarchy problem
- Gauge coupling unification
- Radiative EWSB, light Higgs
- Cold dark matter candidate
- Very rich collider phenomenology

⇒ Most popular and best-studied BSM theory

“Provided superpartners exist at or around the TeV scale”






$$\begin{aligned}\delta m_H^2 &= \left(\frac{g_f^2}{16\pi^2} \right) (\Lambda^2 + m_f^2) - \left(\frac{g_S^2}{16\pi^2} \right) (\Lambda^2 + m_S^2) \\ &= \mathcal{O}\left(\frac{\alpha}{4\pi}\right) |m_S^2 - m_f^2|\end{aligned}$$




Before LHC turn-on

Very optimistic view: if SUSY is light (as we expect...!) it will be discovered early on.





Much easier than discovering the Higgs...

Now



<https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSUS>



Looks grim :- (?



LHC results put supersymmetry theory 'on the spot'




By Pallab Ghosh

Science correspondent, BBC News

Results from the Large Hadron Collider (LHC) have all but killed the simplest version of an enticing theory of sub-atomic physics.

Researchers failed to find evidence of so-called "supersymmetric" particles, which many physicists had hoped would plug holes in the current theory.

Theorists working in the field have told BBC News that they may have to come up with a completely new idea.



Supersymmetry predicts the existence of mysterious super particles.


This looks grim!

Let's reconsider

Arguments for weak-scale SUSY



- Solution to the gauge hierarchy problem
Needs light stops, light higgsinos, somewhat light gluino
- Gauge coupling unification
TeV-scale fermionic states → could be split SUSY
- Radiative EWSB, light Higgs
heavy top effect
 $m_h > 115$ GeV prefers heavy stops (finetuning prize of LEP)
electroweak precision measurements prefer heavy SUSY
- Cold dark matter candidate
TeV-scale LSP could do the job, just needs some efficient annihilation mechanism, e.g. higgsino LSP
- Very rich collider phenomenology
Well, this entirely depends on phase-space ...
and for the time being we are just running at 1/2 force

⇒ Still great, but more “honest” prospects




$$\delta m_H^2 = \left(\frac{g_f^2}{16\pi^2} \right) (\Lambda^2 + m_f^2) - \left(\frac{g_S^2}{16\pi^2} \right) (\Lambda^2 + m_S^2)$$

$$= \mathcal{O}\left(\frac{\alpha}{4\pi}\right) |m_S^2 - m_f^2|$$



Interplay with dark matter searches

Fit to the recent Xenon100 data and its implications for Dark Matter scenarios, taking into account Tevatron and LHC (1.1fb^{-1}) limits



Farina et al., arXiv:1104.3572v3



Beyond CMSSM limits: simplified model spectra

A simplified model is defined by an effective Lagrangian describing the interactions of a small number of new particles. Simplified models can equally well be described by a small number of masses and cross-sections. These parameters are directly related to collider physics observables, making simplified models a particularly [effective framework for evaluating searches](#) and a [useful starting point for characterizing positive signals of new physics](#).

D.Alves et al., arXiv:1105.2838

ATLAS jets+MET search

– interpretation in simplified models: **squarks** –





Exclusion limits in the squark-LSP mass plane for direct [left] and one-step [right] squark decays.

NB: In the case of direct decays, the cross-section for squark_{L,R} production is assumed, however, squark_R production is neglected for the one-step cascade grids, effectively halving the production cross-section. This only applies to the limit contours. For the one-step cascade grids, the nominal cross-sections are too low for any model points to be excluded at 95% C.L., hence no limit contours are drawn.


ATLAS jets+MET search

– interpretation in simplified models: **gluino** –



Exclusion limits in the gluino-LSP mass plane for direct and one-step gluino decays.

CMS results for simplified model spectra



For limits on $m(\tilde{g}), m(\tilde{q}) \gg m(\tilde{g})$ (and vice versa). $\sigma^{\text{prod}} = \sigma^{\text{NLO-QCD}}$.

$$m(\tilde{\chi}^\pm), m(\tilde{\chi}_2^0) \equiv \frac{m(\tilde{g}) + m(\tilde{\chi}^0)}{2}.$$

$m(\tilde{\chi}^0)$ is varied from 0 GeV/c^2 (dark blue) to $m(\tilde{g}) - 200$ GeV/c^2 (light blue).

<https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSUS>

So what does this imply for SUSY in general?

Interpreting LHC SUSY searches in the phenomenological MSSM

S. Sekmen, SK, J. Lykken, S. Moortgat, S. Padhi, L. Pape,
M. Pierini, H.B. Prosper, M. Spiropulu

- We **interpret** within the phenomenological MSSM (pMSSM) the results of **SUSY searches published by the CMS collaboration based on the first 1 fb-1** of data taken during the 2011 LHC run at 7 TeV.
- The pMSSM is a **19-dimensional parametrization of the MSSM** that captures most of its phenomenological features. It encompasses and goes beyond, a broad range of more constrained SUSY models.
- Performing a **global Bayesian analysis**, we obtain **posterior probability densities of parameters, masses and derived observables**.
- In contrast to constraints derived for particular SUSY breaking schemes, such as the CMSSM, our results provide **more generic conclusions on how the current data constrain the MSSM**.

arXiv:1109.5119

pMSSM

- 19-dimensional parametrization of the R-parity conserving MSSM with parameters defined at the SUSY scale
 - the gaugino mass parameters M_1, M_2, M_3 ;
 - the ratio of the Higgs VEVs $\tan \beta = v_2/v_1$;
 - the higgsino mass parameter μ and the pseudo-scalar Higgs mass m_A ;
 - 10 sfermion mass parameters $m_{\tilde{F}}$, where $\tilde{F} = \tilde{Q}_1, \tilde{U}_1, \tilde{D}_1, \tilde{L}_1, \tilde{E}_1, \tilde{Q}_3, \tilde{U}_3, \tilde{D}_3, \tilde{L}_3, \tilde{E}_3$ (imposing $m_{\tilde{Q}_1} \equiv m_{\tilde{Q}_2}, m_{\tilde{L}_1} \equiv m_{\tilde{L}_2}$, etc.),
 - 3 trilinear couplings A_t, A_b and A_τ ,

Assumptions: no new CP phases, flavor-diagonal sfermion mass matrices and trilinear couplings, 1st/2nd generation degenerate and A-terms negligible, lightest neutralino is the LSP.

- Pioneering studies:

“SUSY without prejudice” [C.F. Berger et al., arXiv:0812.0980]

“Fitting the phenomenological MSSM” [S.S. AbdusSalam et al., arXiv:0904.2548]

“The pMSSM leads to a much broader set of predictions for the properties of the SUSY partners as well as for a number of experimental observables than those found in any of the conventional SUSY breaking scenarios such as mSUGRA [CMSSM]. This set of models can easily lead to atypical expectations for SUSY signals at the LHC.”

from the conclusions of arXiv:0812.0980

Method

- Sample the pMSSM parameter space by a Markov-Chain Monte Carlo (MCMC) technique which through a likelihood function incorporates various pre-LHC measurements ($b \rightarrow s\gamma$, $g-2$, ...).
- For a random subset of 500K points, simulate 10K events per point and calculate the signal yields for 3 disjoint CMS SUSY analyses for $\sim 1\text{fb}^{-1}$ of data (α_T hadronic, same-sign dilepton, opposite-sign dilepton)
- Use Bayesian statistics with flat prior to obtain posterior probability distributions for masses, parameters, etc.
- In practice: re-weight the pre-LHC likelihood of these 500K points with the “CMS likelihood” (since the analyses are disjoint, the total likelihood is the product of the individual L’s).
- The main result are posterior distributions.

Tools

- SUSY spectrum calculation: **SOFTSUSY 3.1**
- Low energy observables: **SuperIso 3.0**
- Relic density, DD cross sections: **micrOMEGAs 2.4**
- SUSY mass limits: **micrOMEGAs 2.4**
- Higgs mass limits: **HiggsBounds 2.0.0**
- Decays: **SUSYHIT (SDECAY 1.3b, HDECAY 3.4)**
- SUSY event generation: **PYTHIA 6.4**
- Interfacing: **SUSY Les Houches Accord**
- Generic detector simulation: **DELPHES 1.9**

PS. needed lots of fixes to make everything work....

Analysis setup

- Markov-Chain Monte Carlo sampling of pMSSM parameter space

i	Observable μ_i	Experimental result D_i	Likelihood function $L(D_i \mu_i)$
1	$BR(b \rightarrow s\gamma)$	$(3.55 \pm 0.34) \times 10^{-4}$	Gaussian
2	$BR(B_s \rightarrow \mu\mu)$	$\leq 4.7 \times 10^{-8}$ *)	$1/(1 + \exp(\frac{\mu_2 - D_2}{0.01 D_2}))$
3	$R(B_u \rightarrow \tau\nu)$	1.66 ± 0.54	Gaussian
4	Δa_μ	$(28.7 \pm 8.0) \times 10^{-10} [e^+e^-]$ $(19.5 \pm 8.3) \times 10^{-10} [\text{taus}]$	Weighted Gaussian average
5	m_t	$173 \pm 1.1 \text{ GeV}$	Gaussian
6	$m_b(m_b)$	$4.19^{+0.18}_{-0.06} \text{ GeV}$	Two-sided Gaussian
7	$\alpha_s(M_Z)$	0.1176 ± 0.002	Gaussian
8	m_h	LEP&Tevatron (HiggsBounds)	$L_8 = 1$ if allowed. $L_8 = 10^{-9}$ if m'_h sampled from $Gauss(m_h, 1.5)$ is excluded.
9	sparticle masses	LEP (micrOMEGAs)	$L_9 = 1$ if allowed $L_9 = 10^{-9}$ if excluded

*) we re-weight a posteriori with the new limit $BR(B_s \rightarrow \mu\mu) < 1.08$ at 95% CL

$$\begin{aligned}
 |M_i| &\leq 3 \text{ TeV} \\
 |\mu| &\leq 3 \text{ TeV} \\
 m_A &\leq 3 \text{ TeV} \\
 m_{\tilde{F}} &\leq 3 \text{ TeV} \\
 |A_{t,b,\tau}| &\leq 7 \text{ TeV} \\
 2 &\leq \tan \beta \leq 60
 \end{aligned}$$

- We use a flat prior for parameters and sample 1.5×10^7 points
- Distribution of points maps the total likelihood, $L_{\text{preLHC}} = \sum_{i=1}^9 L_i$
- Draw a random subset of 5×10^5 points for simulation



We consider the following 3 public disjoint CMS SUSY analyses using $\sim 1 \text{ fb}^{-1}$ 7 TeV 2011 data:

- **Hadronic α_T** (RA1 – CMS-SUS-11-003) 1.1 fb^{-1} : ≥ 2 jets and $\alpha_T > 0.55$, where α_T is designed to distinguish between real and fake E_T^{miss} . **Shape analysis** that gives results in 8 disjoint H_T bins. We use **all**.
- **Same sign (SS) dilepton** (RA5 - CMS-SUS-11-010) 0.98 fb^{-1} : **Opposite sign dilepton** analysis. 8 correlated analysis regions. We use $H_T > 400$, $E_T^{\text{miss}} > 120$.
- **Opposite sign (OS) dilepton** (RA6 – CMS-SUS-11-011) 0.98 fb^{-1} : **Same sign dilepton analysis**. 2 correlated analysis regions. We use $H_T > 300$, $E_T^{\text{miss}} > 275$.

Slide from Sezen's talk, 30-08-11 at LHC2TeV workshop, CERN

CMS results used

j	Analysis and search region (values in GeV)	Observed event count (N_j)	Data-driven SM BG estimate ($B_j \pm \delta B_j$)
1	α_T hadronic, $275 \leq H_T < 325$	782	$787.4^{+31.5}_{-22.3}$
2	α_T hadronic, $325 \leq H_T < 375$	321	$310.4^{+8.4}_{-12.4}$
3	α_T hadronic, $375 \leq H_T < 475$	196	$202.1^{+8.6}_{-9.4}$
4	α_T hadronic, $475 \leq H_T < 575$	62	$60.4^{+4.2}_{-3.0}$
5	α_T hadronic, $575 \leq H_T < 675$	21	$20.3^{+1.8}_{-1.1}$
6	α_T hadronic, $675 \leq H_T < 775$	6	$7.7^{+0.8}_{-0.5}$
7	α_T hadronic, $775 \leq H_T < 875$	3	$3.2^{+0.4}_{-0.2}$
8	α_T hadronic, $875 \leq H_T$	1	$2.8^{+0.4}_{-0.2}$
9	SS 2ℓ , $H_T > 400$, $\cancel{E}_T > 120$	1	2.3 ± 1.2
10	OS 2ℓ , $H_T > 300$, $\cancel{E}_T > 275$	8	4.2 ± 1.3

Assume a Poisson likelihood $\text{Poisson}(N_j | s_j + b_j)$ with expected count $s_j + b_j$ and compute the marginal likelihood $L_j = p(N_j | s_j)$ by integrating over the expected background b_j



Likelihood:


$$\begin{aligned}
 L_{CMS} &\equiv p(N|s, b, \delta b) \\
 &= \prod_{i=1}^{10} \text{Poisson}(N_i|s_i + b_i) p(b_i|\delta b_i) && \text{takes into account the} \\
 & && \text{uncertainty in the BG} \\
 & && \text{estimate} \\
 &= \prod_{i=1}^{10} \frac{e^{-(s_i+b_i)} (s_i + b_i)^{N_i}}{N_i!} \frac{e^{-k_i b_i} (k_i b_i)^{Q_i}}{\Gamma(Q_i + 1)}
 \end{aligned}$$


where $Q_i = (B_i/\delta B_i)^2$

$$k_i = B_i/\delta B_i^2$$




Slide from Sezen's talk, 30-08-11 at LHC2TeV workshop, CERN

Results: posterior densities








BCR: Bayesian credible region – these are not exclusion curves!




neutralino
vs. gluino




chargino
vs. gluino

BCR: Bayesian credible region — these are not exclusion curves!



Dark matter implications




$$p(\Omega h^2 < 0.136) = 55\%$$


$$p(0.094 < \Omega h^2 < 0.136) = 1\%$$

Implications for dark matter searches

spin-independent




spin-dependent



$$\xi = \Omega h^2 / 0.1123$$

Implications for Higgs



Complementary study

“Implications of LHC Searches on SUSY Particle Spectra: The pMSSM Phase Space with Neutralino Dark Matter”



A. Arbey, M. Battaglia, F. Mahmoudi

- Study the implications of LHC searches on SUSY particle spectra using **flat scans** of the 19-parameter pMSSM phase space.
- **Apply 2σ constraints** from flavour physics, $g-2$, dark matter and earlier LEP and Tevatron searches.
- The sensitivity of the LHC SUSY searches with jets, leptons and missing energy is assessed by **reproducing with fast simulation the recent CMS analyses** after validation on benchmark points.
- Present results in terms of the **fraction of pMSSM points** compatible with all the constraints **which are excluded by the LHC searches with 1 fb^{-1} and 15 fb^{-1}** as a function of the mass of strongly and weakly interacting SUSY particles.
- Discuss the **suppression of Higgs production cross sections** for the MSSM points not excluded and **contrast the region of phase space tested by the LHC data with the constraints from dark matter direct detection experiments.**

arXiv:1110.3726

- Fraction of accepted points from flat random scan

arXiv:1110.3726





$$2.16 \times 10^{-4} < \text{BR}(B \rightarrow X_s \gamma) < 4.93 \times 10^{-4}$$

$$\text{BR}(B_s \rightarrow \mu^+ \mu^-) < 1.26 \times 10^{-8}$$





$$-2.4 \times 10^{-9} < \delta a_\mu < 4.5 \times 10^{-9}$$

$$10^{-4} < \Omega_\chi h^2 < 0.135$$





- Fraction of accepted points excluded by CMS

arXiv:1110.3726



- Interplay with direct dark matter detection


arXiv:1110.3726



black dots: accepted pMSSM points,
 grey dots: points not excluded by 1 fb^{-1} and the
 light grey dots: points not excluded by 15 fb^{-1} of LHC data.

- Prediction for light Higgs searches

arXiv:1110.3726







$$R = \frac{\sigma(gg \rightarrow h^0)}{\sigma(gg \rightarrow H_{SM}^0)} \times \frac{\text{BR}(h^0 \rightarrow \gamma\gamma)}{\text{BR}(H_{SM}^0 \rightarrow \gamma\gamma)}$$

Natural SUSY endures

M. Papucci, J.T. Ruderman, A. Weiler

From the requirement of naturalness

- stops & left sbottom below 500–700 GeV
- higgsinos below 200–350 GeV
- a not too heavy gluino, 900–1500 GeV



Conclusions

- LHC results are pushing squark and gluino mass limits to ~ 1 TeV; expectations for early discoveries were too optimistic
- Current searches are not (yet) sensitive to
 - ★ Small mass differences $\rightarrow$ soft jets, low E_T^{miss}
 - ★ Compressed spectra in general
 - ★ Mainly electroweak production
 - ★ Mainly stop/sbottom production
- Plenty of room where SUSY can hide
besides, EW fits and flavor physics actually prefer heavy SUSY
- SUSY DM stays compelling case
interesting complementarity between LHC and DD
- We definitely need [the means] to interpret LHC results in terms of a wide range of models, including pMSSM.



Needs communication between experimentalists (collaborations!) and theorists

Implications of LHC results for TeV-scale physics

from 29 August 2011 to 02 September 2011 (Europe/Zurich)

Europe/Zurich timezone

Overview

Timetable

Registration

Registration Form

List of registrants

Video Services

Arrival Information

This Workshop, which takes place during the last week of the 2011 TH/LPCC Institute on LHC physics, will be the first in a series of meetings devoted to evaluating the implications of recent results from the LHC, and elsewhere, for TeV-scale physics, and to discuss the impact of these results on the future strategy for particle physics.

The Workshop continues the activity of the 2009 TH Institute "From the LHC to a Future Collider", <https://indico.cern.ch/conferenceDisplay.py?confId=40437>, whose findings were documented in <http://arXiv.org/abs/0909.3240>.

The results will be summarised in a document, to be submitted as input to the planned 2012 update of the European Strategy for Particle Physics. The participation and input of all those interested, experimentalists and theorists, is welcome and encouraged!

The workshop activities will be focused around three working groups investigating different aspects of TeV-scale physics, namely


- Signals of electroweak symmetry breaking (conveners: S.Heinemyer, M.Kado, C.Mariotti, G.Weiglein, A.Weiler)
- Signatures with missing energy (conveners: R.Cavenaugh, J.Hewett, S.Kraml, G.Polesello)
- Other signatures of possible BSM physics (conveners: C.Grojean, D.Martinez, J.Santiago Perez, P.Savard, S.Worm)

The task of the working groups is to assess the possible interpretations of the experimental results in view of their implications for the future strategy of particle physics.

The charge for the first meeting is to summarise the experimental situation at this time, to start the discussion of possible interpretations, and to define the lines of work that should be carried out. We expect this to be followed by more informal meetings of the working groups, leading to a final plenary meeting, foreseen to take place in May or June 2012, such that the final document will be ready in time for the 'Orsay-type' meeting of the European Strategy update. The precise dates will be decided once the timeline of the Strategy process is finalized by CERN Council.

Next plenary meeting after Moriond (week of March 26) at CERN

ATLAS leptons+MET



Requiring $\Omega h^2 < 0.136$

