
Exercises String Theory
Prof. Dr. Albrecht Klemm

1 The propagator of the free boson

2 points

In two dimensions, the free boson has the following Euclidean action

$$S = \frac{1}{2}g \int d^2x (\partial_\mu \phi \partial^\mu \phi + m^2 \phi^2). \quad (1.1)$$

1. Show that the propagator

$$K(x, y) = \langle \phi(x) \phi(y) \rangle \quad (1.2)$$

obeys

$$g(-\partial_x^2 + m^2)K(x, y) = \delta(x - y). \quad (1.3)$$

2. Show that $K(x, y)$ only depends on $r = |x - y|$ and that

$$K(r) = -\frac{1}{2\pi g} \ln r. \quad (1.4)$$

Hint: Integrate (1.3) and derive an ordinary differential equation for $K(r)$.

3. Rewrite $K(r)$ in complex coordinates. Show that

$$\begin{aligned} \langle \partial_z \phi(z) \phi(w) \rangle &= -\frac{1}{4\pi g} \frac{1}{z - w} \\ \langle \partial_z \phi(z) \partial_w \phi(w) \rangle &= -\frac{1}{4\pi g} \frac{1}{(z - w)^2} \end{aligned} \quad (1.5)$$

2 The ghost system

4 points

The action for the ghost system is given by

$$S = \frac{1}{2} \int d^2x b_{\mu\nu} \partial^\mu c^\nu. \quad (2.1)$$

Both, c and $b_{\mu\nu}$ are anti-commuting and $b_{\mu\nu}$ is a traceless tensor. The propagator is found to be

$$\langle b(z)c(w) \rangle = \frac{1}{\pi g} \frac{1}{z - w}. \quad (2.2)$$

1. Determine the correlators

$$\langle b(z) \partial_w c(w) \rangle, \quad \langle \partial_z b(z) c(w) \rangle \quad \text{and} \quad \langle \partial_z b(z) \partial_w c(w) \rangle. \quad (2.3)$$

The normal ordered energy-momentum tensor of the system is given by

$$T(z) = \pi g : (2\partial cb + c\partial b) : \quad (2.4)$$

2. Compute the OPE's $T(z)c(w)$ and $T(z)b(w)$ using Wick's theorem and read of the corresponding conformal dimensions.
3. Compute the OPE of T with itself

$$T(z)T(w) = \frac{c/2}{(z-w)^4} + \frac{hT(w)}{(z-w)^2} + \frac{\partial T(w)}{(z-w)} \quad (2.5)$$

and determine the central charge as well as the conformal weight.

3 Vertex operators

4 points

Bosonic Vertex operators are defined by

$$V_\alpha(z, \bar{z}) =: e^{i\alpha\phi(z, \bar{z})} : \quad (3.1)$$

where ϕ is a free bosonic field.

1. Show that the OPE of $\partial\phi$ with V_α is

$$\partial\phi(z)V_\alpha(w, \bar{w}) = -\frac{i\alpha}{4\pi g} \frac{V_\alpha(w, \bar{w})}{z-w} + \dots \quad (3.2)$$

2. Show that the OPE of V_α with T is given as

$$T(z)V_\alpha(w, \bar{w}) = \frac{\alpha^2}{8\pi g} \frac{V_\alpha(w, \bar{w})}{(z-w)^2} + \frac{\partial_w V_\alpha(w, \bar{w})}{z-w} + \dots \quad (3.3)$$

What is the conformal dimension?

3. Show that for two free fields one has

$$: e^{a\phi_1} :: e^{b\phi_2} :=: e^{a\phi_1 + b\phi_2} : e^{ab\langle\phi_1\phi_2\rangle}. \quad (3.4)$$

Use this result to compute the OPE of two Vertex operators.