

Exercises String Theory
 Prof. Dr. Albrecht Klemm

1 The conformal and the Lorentz group

4 points

We recall that the generators of the conformal group can be expressed as follows

translation	$P_\mu = -i\partial_\mu$
dilatation	$D = -ix^\mu\partial_\mu$
rotation	$L_{\mu\nu} = i(x_\mu\partial_\nu - x_\nu\partial_\mu)$
special conformal transformation	$K_\mu = -(2x_\mu x^\nu\partial_\nu - x^2\partial_\mu)$

They obey the following algebra

$$\begin{aligned}
 [D, P_\mu] &= iP_\mu, & [D, K_\mu] &= -iK_\mu, & [K_\mu, P_\nu] &= 2i(\eta_{\mu\nu}D - L_{\mu\nu}), \\
 [K_\rho, L_{\mu\nu}] &= i(\eta_{\rho\mu}K_\nu - \eta_{\rho\nu}K_\mu), & [P_\rho, L_{\mu\nu}] &= i(\eta_{\rho\mu}P_\nu - \eta_{\rho\nu}P_\mu), \\
 [L_{\mu\nu}, L_{\rho\sigma}] &= i(\eta_{\nu\rho}L_{\mu\sigma} + \eta_{\mu\sigma}L_{\nu\rho} - \eta_{\mu\rho}L_{\nu\sigma} - \eta_{\nu\sigma}L_{\mu\rho})
 \end{aligned} \tag{1.1}$$

1. Check the first four relations.
2. Show that the re-defined fields

$$\begin{aligned}
 J_{\mu\nu} &= L_{\mu\nu} & J_{-1,\mu} &= \frac{1}{2}(P_\mu - K_\mu) \\
 J_{-1,0} &= D & J_{0,\mu} &= \frac{1}{2}(P_\mu + K_\mu)
 \end{aligned} \tag{1.2}$$

obey the $SO(d+1,1)$ commutation relations

$$[J_{ab}, J_{cd}] = i(\eta_{ad}J_{bc} + \eta_{bc}J_{ad} - \eta_{ac}J_{bd} - \eta_{bd}J_{ac}). \tag{1.3}$$

2 Two-point function

2 points

We consider the two-point function of two primary fields Φ_i , with conformal weight Δ_i , $i = 1, 2$ in 2d CFT

$$G(z_1, z_2) = \langle \Phi_1(z_1)\Phi_2(z_2) \rangle. \tag{2.1}$$

Recall that under infinitesimal transformation $z \rightarrow z + \epsilon$ the fields transform as

$$\delta_\epsilon \Phi_i(z) = (\epsilon(z)\partial + \Delta_i\partial\epsilon(z))\Phi_i, \quad i = 1, 2. \tag{2.2}$$

1. Show that conformal invariance implies

$$(\epsilon(z_1)\partial_1 + \Delta_1\partial\epsilon(z_1) + \epsilon(z_2)\partial_2 + \Delta_2\partial\epsilon(z_2))G(z_1, z_2) = 0. \tag{2.3}$$

2. By setting $\epsilon = 1$, show that $G(z_1, z_2)$ is a function of $x = z_1 - z_2$ only.
3. By setting $\epsilon = z$, show that $G(z_1, z_2)$ takes the following form

$$G(x) = \frac{C}{x^{\Delta_1 + \Delta_2}}, \quad (2.4)$$

where C is a constant.

4. By setting $\epsilon = z^2$, show that $G(z_1, z_2)$ vanishes unless $\Delta_1 = \Delta_2$.

3 Global conformal transformations

2 points

As is known from the lecture, the invariance under global conformal transformations can be encoded in the following differential equations imposed on the correlators of primary fields

$$\begin{aligned} \sum_i \partial_{w_i} \langle \phi_1(w_1) \dots \phi_n(w_n) \rangle &= 0 \\ \sum_i (w_i \partial_{w_i} + \Delta_i) \langle \phi_1(w_1) \dots \phi_n(w_n) \rangle &= 0 \\ \sum_i (w_i^2 \partial_{w_i} + w_i \Delta_i) \langle \phi_1(w_1) \dots \phi_n(w_n) \rangle &= 0. \end{aligned} \quad (3.1)$$

Show explicitly for the two- and the three-point function that these relations are indeed valid.

4 Operator product expansion

2 points

Starting with

$$\delta_{\epsilon, \bar{\epsilon}} \phi(z, \bar{z}) = [Q_\epsilon + Q_{\bar{\epsilon}}, \phi(z, \bar{z})], \quad Q_\epsilon = \oint_{C_0} \frac{dz}{2\pi i} \epsilon(z) T(z) \quad (4.1)$$

and

$$\delta_{\epsilon, \bar{\epsilon}} \phi(z, \bar{z}) = (\Delta \partial \epsilon + \bar{\Delta} \bar{\partial} \epsilon + \epsilon \partial + \bar{\epsilon} \bar{\partial}) \phi(z, \bar{z}), \quad (4.2)$$

Show that

$$T(z) \phi(w, \bar{w}) = \frac{\Delta}{(z-w)^2} \phi(w, \bar{w}) + \frac{1}{z-w} \partial_w \phi(w, \bar{w}) + \text{reg}. \quad (4.3)$$

In the last expression, radial ordering is understood. Hint: Use the operator product expansion

$$\phi_i(z, \bar{z}) \phi_j(w, \bar{w}) = \sum_k c_{ijk} (z-w)^{\Delta_k - \Delta_i - \Delta_j} (\bar{z}-\bar{w})^{\bar{\Delta}_k - \bar{\Delta}_i - \bar{\Delta}_j} \quad (4.4)$$

and the Cauchy formula

$$\oint_{C_w} \frac{dz}{2\pi i} \frac{f(z)}{(z-w)^n} = \frac{1}{(n-1)!} f^{(n-1)}(w) \quad (4.5)$$