

Condensed Matter Theory I — WS09/10

Exercise 6

(Please return your solutions before Fr., 15.1., 12:00h)

6.1 Phonons in a metal: Kohn Anomaly (10 points)

In metals one expects that the harmonic interactions between the ions of a lattice are not only between nearest neighbors, but are long-ranged. The physical origin is a displacement of an ionic charge that induces an electric charge polarization in the electron sea which is long-ranged and oscillatory in space. This oscillatory polarization acts on the other ions in the lattice. The oscillations are induced by the fact that the electronic states are filled up to the Fermi wave number k_F and the wave number of the resulting oscillatory ion-ion potential is $2k_F$ (Friedel oscillations).

We now consider a one-dimensional linear chain of ions in an electron sea with the lattice constant a . The force constant between an ion on lattice site i and an ion on lattice site j is

$$\kappa_{ij} = \kappa_0 \frac{\sin 2k_F a(i-j)}{k_F a(i-j)}.$$


- Write down the Lagrange function for the system of ions and derive the equations of motion.
- Make an ansatz of plane wave solutions for the ion displacements:

$$q_j(t) = q_0 e^{i(kx - \omega t)}, \quad x_j = j \cdot a$$

Derive an expression for $\omega(k)^2$ and $\frac{\partial \omega^2}{\partial k}$ and show that $\omega(k)^2$ has a cusp (divergent slope) at $k = 2k_F$. This effect is called Kohn anomaly after Walter Kohn, who predicted it in 1959.

6.2 Optical phonons (10 points)

Consider a one-dimensional harmonic chain with two different atoms in the primitive cell,



where $x_j = a \cdot j$ is the position of the j th primitive cell and q_{sj} is the elongation of the s th atom in the j th cell.

The Hamiltonian of the system is given by

$$H = \sum_{js} \frac{1}{2} M_s \dot{q}_s(x_j) + \frac{1}{2} \sum_j \kappa_{12} (q_{2j} - q_{1j})^2 + \kappa_{21} (q_{1j} - q_{2j-1})^2.$$

For simplicity we assume $\kappa_{12} = \kappa_{21} = \kappa$.

- (a) Write down the equations of motion. Use the ansatz

$$q_{sj}(t) = q_s^0 e^{i(ka_j - \omega t)}, \quad s = 1, 2$$

to derive the following equation for the frequencies of the eigenmodes (phonons):

$$\omega^2 = \frac{\kappa}{M^*} \pm \sqrt{\left(\frac{\kappa}{M^*}\right)^2 - 2\frac{\kappa^2}{M_1 M_2}(1 - \cos ka)}$$

$M^* = \frac{M_1 M_2}{M_1 + M_2}$ is the reduced mass.

- (b) Calculate $\omega(k)^2$ for $k \rightarrow 0$ and $k \rightarrow \pm \frac{\pi}{a}$ and sketch its behavior for the full Brillouin zone $[-\frac{\pi}{a}, \frac{\pi}{a}]$. Identify the acoustic and the optical branch.
- (c) Discuss the limit $M_1 = M_2$ and show that this limit results in the acoustic phonon dispersion with lattice constant $\frac{a}{2}$.

6.3 Meissner effect

(10 points)

The London equations of superconductivity are


$$\frac{\partial}{\partial t} (\Lambda \vec{j}) = \vec{E} \quad \vec{\nabla} \times (\Lambda \vec{j}) = -\vec{H} \quad \Lambda = \frac{m}{2n_s e^2}.$$

Consider a thin superconducting slab of thickness d , infinitely extended in y and z direction, in a uniform, static magnetic field $\vec{H}_0 = H_0 \hat{e}_z$ parallel to the slab surface.

- a) Use the Maxwell equations to calculate the magnetic field $\vec{H}(\vec{x})$ inside the slab. Show that the field enters the slab only on a length scale λ_L (London penetration depth). Draw the result.
- b) Calculate also the current density inside the slab and draw the result.
- c) Calculate the average magnetization of the slab,

$$\langle H(x) \rangle = \frac{1}{d} \int_{-d/2}^{d/2} dx H(x),$$

and draw it as a function of d .




Consider now a current flow through an infinitely long cylindrical superconducting wire of radius R ($R \gg \lambda_L$).

- d) Use the continuity equation to show that the current density distribution $\vec{j}(r)$ obeys an ordinary differential equation of the form (*modified Bessel differential equation*)

$$x^2 \frac{d^2 j}{dx^2} + x \frac{dj}{dx} - x^2 j = 0$$

Hint: Laplace operator in cylindrical coordinates

$$\Delta = \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \phi^2} + \frac{\partial^2}{\partial z^2}$$



- e) Draw the solution of d) (*modified Bessel function of the first kind*) and show that the current flows only in a small layer underneath the surface of the wire.
- f) Use Biot-Savart's law to calculate the magnetic field outside the wire induced by the current. If the magnetic field at the surface exceeds a critical value H_c , the superconductivity will collapse. What is the corresponding critical current I_c ?