

## Übungen zur Festkörpertheorie I — WS03/04

### 4. Übungsblatt

#### 1. Band structure of a 2D surface band

Above the surface of a metal there often exists a surface band of electrons, i.e. a band of electrons moving in strictly two-dimensional (2D) space parallel to the surface. The precise origin of a surface band will be discussed later in the lecture. Here we want to consider the band structure  $\varepsilon_k$ .

On a plain surface, the 2D electrons move almost freely (mass  $m$ ), since the surface band states are somewhat above the surface and, hence, are decoupled from the periodic potential of the lattice.

On gold surfaces, there, in addition, is a lattice reconstruction (so-called herring bone reconstruction, Fig. 1) where the surface is modulated in the  $x$  direction with the periodic potential shown in Fig. 2,

$$U(x) = U_0 \Theta(\cos(\pi x/a)) , \quad (1)$$

while the  $y$  direction remains translationally invariant.

- a) Calculate the 2D surface band structure  $\varepsilon_k$ . Distinguish energies  $E > U_0$  and  $E < U_0$ .
- b) Sketch the lines of constant energy on the surface.

#### 2. Electron motion in a herring bone surface band

We describe the surface band approximately by the band structure

$$\varepsilon_k = D[1 - \cos(2ak_x)] + \frac{(\hbar k_y)^2}{2m} \quad (2)$$

and apply an electric field  $\vec{E} = (E_x, E_y) = \text{const.}$ .

Calculate the quasiclassical motion of an electron in this band structure. Sketch the trajectories in the  $x$ - $y$  plane.




Figure 1: Herring bone reconstruction




Figure 2: Periodic potential