

---

## Exercises on Elementary Particle Physics

Prof. Dr. H.-P. Nilles

### 1. *Non-Abelian Gauge Symmetry - part III*

In Exercise 5, we discussed non-abelian gauge symmetries. We showed that the Lagrangian

$$\mathcal{L} = \bar{\Psi}(x) (i\gamma^\mu D_\mu) \Psi(x) - \frac{1}{2} \text{tr} (F_{\mu\nu} F^{\mu\nu})$$

is invariant under local gauge transformations, where  $F_{\mu\nu} = F_{\mu\nu}^a T^a$  and the generators of the algebra  $T^a$  are normalized to  $\text{tr}(T^a T^b) = \frac{1}{2} \delta^{ab}$ .

- (a) Show that, in the case of SU(2), the normalization condition is fulfilled by the Pauli-matrices  $T^a = \frac{1}{2} \sigma^a$ .
- (b) Prove the following equation:

$$\frac{1}{2} \text{tr} (F_{\mu\nu} F^{\mu\nu}) = \frac{1}{4} F_{\mu\nu}^a F^{\mu\nu a}$$

## 2. The Standard Model Higgs Effect - part I

In the Standard model, the electroweak interactions of leptons are described by the Lagrangian

$$\begin{aligned}
 \mathcal{L} = & \bar{R}i\gamma^\mu D_\mu R + \bar{L}i\gamma^\mu D_\mu L && \text{kinetic energy of leptons and interactions} \\
 & -\frac{1}{4}G_{\mu\nu}^a G^{\mu\nu a} - \frac{1}{4}F_{\mu\nu} F^{\mu\nu} && \text{with gauge bosons} \\
 & + (D_\mu \phi)^\dagger (D^\mu \phi) - \mu^2 \phi^\dagger \phi - \lambda (\phi^\dagger \phi)^2 && \text{kinetic energy of gauge bosons and} \\
 & -G_e (\bar{L}\phi R + \bar{R}\phi^\dagger L) && \text{self-interactions} \\
 & && \text{Higgs field with potential} \\
 & && \text{electron mass and coupling to Higgs}
 \end{aligned}$$

with the covariant derivative

$$D_\mu = \partial_\mu + ig' \frac{Y}{2} B_\mu + ig T^a W_\mu^a$$

and the particle content

|   | Hypercharge $Y$ | rep. of $SU(2)_L$ | rep. of Lorentz algebra      |
|---|-----------------|-------------------|------------------------------|
| $L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$        | -1              | <b>2</b>          | $(\frac{1}{2}, 0)$ (*)       |
| $R = e_R$   | -2              | <b>1</b>          | $(0, \frac{1}{2})$ (*)       |
| $\phi = \begin{pmatrix} \phi_+ \\ \phi_0 \end{pmatrix}$ | 1               | <b>2</b>          | $(0, 0)$                     |
| $T^a W_\mu^a$   | 0               | <b>3</b>          | $(\frac{1}{2}, \frac{1}{2})$ |
| $B_\mu$   | 0               | <b>1</b>          | $(\frac{1}{2}, \frac{1}{2})$ |

(\*) for now, L and R contain Dirac spinors

- Write down how the covariant derivative acts on the left-handed lepton doublet, on the right-handed lepton and on the Higgs doublet.
- Show that the Lagrangian is Lorentz invariant.
- Show that the Lagrangian is gauge invariant.
- Find the minimum of the Higgs potential for  $\mu^2 < 0$  and choose the vacuum expectation value (vev) to be

$$\langle \Phi \rangle_0 := \langle 0 | \Phi | 0 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}.$$

- (e) We want to reparametrize the shifted Higgs field (which contains two complex fields  $\phi_+$  and  $\phi_0$ ) in terms of the four real fields  $\xi^a$  ( $a = 1, 2, 3$ ) and  $\eta$  in the following way:

$$\phi(x) = \phi'(x) + \langle \phi \rangle_0 = U^{-1}(x) \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}}(v + \eta(x)) \end{pmatrix}, \quad U(x) = \exp \left( -\frac{i}{v} \xi^a(x) T^a \right)$$

After this redefinition, we apply a  $SU(2)_L$  gauge transformation to all  $SU(2)_L$  non-singlets ( $L, \phi$  and  $T^a W_\mu^a$ ). This gauge is called the unitary gauge. In this gauge, the Higgs field has the simple form:

$$\phi \mapsto \phi' = U(x)\phi = \begin{pmatrix} 0 \\ \frac{1}{\sqrt{2}}(v + \eta(x)) \end{pmatrix}$$

In the following, we skip the prime for all fields. Show that the Higgs potential in this gauge now reads

$$V(\phi) = \mu^2 \eta^2 + \lambda v \eta^3 + \frac{\lambda}{4} \eta^4.$$

What is the mass of the  $\eta$  field? Compare the degrees of freedom in the Higgs sector to the situation before symmetry breakdown.

- (f) Next, we consider the kinetic energy of the Higgs field. Show that:

$$\begin{aligned} (D_\mu \phi)^\dagger (D^\mu \phi) &= \frac{1}{2} \partial_\mu \eta \partial^\mu \eta + \frac{1}{4} g^2 (v + \eta)^2 W_\mu^- W^{\mu+} \\ &\quad + \frac{1}{8} (v + \eta)^2 (W_\mu^3, B_\mu) \begin{pmatrix} g'^2 & -g'g \\ -g'g & g^2 \end{pmatrix} \begin{pmatrix} W^{\mu 3} \\ B^\mu \end{pmatrix} \end{aligned}$$

where the matrix will lead to the mass matrix in the following.

- (g) The masses are given by the terms that are quadratic in the fields, e.g.

$$\frac{1}{4} g^2 v^2 W_\mu^- W^{\mu+} = m_W^2 W_\mu^- W^{\mu+}$$

so  $m_W = \frac{1}{2} v g$ . Obviously, this is not so easy for  $W_\mu^3$  and  $B_\mu$ . To see the masses of these fields we have to diagonalize the mass matrix

$$\frac{1}{8} v^2 (W_\mu^3, B_\mu) O^T O \begin{pmatrix} g'^2 & -g'g \\ -g'g & g^2 \end{pmatrix} O^T O \begin{pmatrix} W^{\mu 3} \\ B^\mu \end{pmatrix} =: \frac{1}{2} (Z_\mu, A_\mu) \begin{pmatrix} m_Z^2 & 0 \\ 0 & m_A^2 \end{pmatrix} \begin{pmatrix} Z^\mu \\ A^\mu \end{pmatrix}$$

with an orthogonal matrix. Determine this orthogonal matrix by computing the eigenvalues and the eigenvectors of the mass matrix.

What are the masses of the  $Z_\mu$  and  $A_\mu$  fields?

Compare the degrees of freedom in the gauge sector to the situation before symmetry breakdown. How about the total amount of degrees of freedom?

- (h) On the other hand, one can always write an orthogonal  $2 \times 2$  matrix in the following way:

$$O = \begin{pmatrix} \cos \theta_W & -\sin \theta_W \\ \sin \theta_W & \cos \theta_W \end{pmatrix}$$

Write  $\cos \theta_W$  in terms of  $g'$  and  $g$ . Prove that the ratio of the masses of the  $W$ - and the  $Z$ -boson is:

$$\frac{m_W}{m_Z} = \cos \theta_W$$

This is a prediction of the standard model, which has been experimentally confirmed within a small error.

- (i) Next, we consider the covariant derivative.

$$D_\mu = \partial_\mu + ig' \frac{Y}{2} B_\mu + ig T^a W_\mu^a$$

Substitute the fields  $B_\mu$  and  $W_\mu^a$  by  $W_\mu^\pm$ ,  $Z_\mu$  and  $A_\mu$ . Show that it follows:

$$D_\mu = \partial_\mu + iA_\mu \frac{g'g}{\sqrt{g'^2 + g^2}} \left( T_3 + \frac{Y}{2} \right) + iZ_\mu \frac{1}{\sqrt{g'^2 + g^2}} \left( g^2 T_3 - g'^2 \frac{Y}{2} \right) + \frac{ig}{\sqrt{2}} \begin{pmatrix} 0 & W_\mu^+ \\ W_\mu^- & 0 \end{pmatrix}$$

Now, we can define the electric charge:

$$e := \frac{g'g}{\sqrt{g'^2 + g^2}} \quad \text{and} \quad Q := T_3 + \frac{Y}{2}$$