

Condensed Matter Theory I — WS05/06

Numerical Exercise

N.1 Hartree-Fock Approximation

In one of the previous exercises we found an expression for the occupation number of spin-up and spin-down electrons in the presence of a local potential within a Hartree-Fock approximation:

$$\langle n_\sigma \rangle = \frac{1}{N} \sum_k \left\{ 1 + \exp \left(\frac{\varepsilon_k + U \langle n_{-\sigma} \rangle - \mu}{T} \right) \right\}^{-1} \quad (1)$$

The absolute value of the magnetization is then given by

$$m = |\langle n_\uparrow \rangle - \langle n_\downarrow \rangle|$$

Assume that the density of states has the shape of a half circle, i.e.

$$N(\varepsilon) = \frac{2N}{\pi} \cdot \sqrt{D^2 - \varepsilon^2}$$

and choose $D = 1$ and $U = 2$.

Solve then the two equations for $\langle n_\uparrow \rangle$ and $\langle n_\downarrow \rangle$ self consistently in the temperature interval $T \in [0, 1]$, i.e. start with non-symmetric distributions (for instance $\langle n_\uparrow \rangle = 0.55$ and $\langle n_\downarrow \rangle = 0.45$), calculate first with these distributions the chemical potential such that the condition

$$\langle n_\uparrow \rangle + \langle n_\downarrow \rangle = 1$$

is fulfilled and then calculate in a second step the right side of (1). The results will define new distributions. Now iterate this procedure until the results will be stable. Plot m as a function of T . (The result should look like the plot in Fig. 1)

Hint: The simplest way to calculate the integral numerically is to use an equidistant grid and to approximate the integral as a discrete sum over the gridpoints.

The chemical potential can be found by using a sequence of nested intervals.

N.2 Coupled Differential Equations

In this exercise we want to try to find numerical solutions for two different systems of coupled differential equations.

- a) The first example is the following well-known system of two coupled differential equations

$$\begin{aligned} f'(x) &= g(x) & \text{with the boundary condition:} & & f(0) &= 0 \\ g'(x) &= -f(x) & & & g(0) &= 1 \end{aligned}$$

Rewrite these equations as integral equations and calculate the two functions in the interval $x \in [0, 2\pi]$ and compare the numerical result with the exact solution.

- b) The second example is a typical example for a differential equation as it appears in *renormalization group* calculations.

$$f'(x) = -\frac{f^2(x)}{x} \quad \text{with: } f(1) = f_0$$

Calculate f for $x \in [0, 1]$ for the two cases $f_0 = \pm 1$.

- c) Consider now the more general case

$$\begin{aligned} f'(x) &= -\frac{g^2(x)}{x} & f(1) &= f_0 \\ g'(x) &= -\frac{f(x) \cdot g(x)}{x} & g(1) &= g_0 \end{aligned} \quad \text{with:}$$

Solve this system for different values of f_0 and g_0 .

Physical background: These equations arise by studying the influence of a single magnetic impurity in a non-magnetic host ('Kondo model'). Its solution contains very important information about the low temperature behaviour of a spin-spin-coupling between the impurity and the band electrons.




Figure 1: The magnetization as a function of temperature